Rickets and Its Associated Factors among Under-Five Children in Selected Public Hospitals in Eastern Ethiopia

Kaleab Sisay¹, Firehiwot Mesfin², Tesfaye Gobena³, Berhe Gebremichael^{4*}

Abstract

Background: Rickets affects the rapid growth of children. It is prevalent in low-income countries, and ranks among the five most common diseases in children. However, its magnitude and associated factors are not well studied and documented in eastern Ethiopia. Therefore, the aim of this study was to assess the magnitude and the associated factors of rickets among under-five children in selected public hospitals of eastern Ethiopia.

Methods: A hospital based cross-sectional study was conducted from February 23 to March 10, 2017 among 590 under-five children in Hiwot Fana Specialized University Hospital, and Haramaya and Dilchora General Hospitals. Consecutive sampling was used to reach the actual respondents. Data were collected using a pre-tested questionnaire and the presence of rickets was assessed based on clinical signs and symptoms. Data were entered into EpiData Version 3.1 and exported into SPSS Version 22.0 for analysis. Bivariable and multivariable logistic regression models were fitted. Level of significance was considered at p-value less than 0.05.

Results: The magnitude of rickets was 7.8% (95% CI: 5.70, 10.20). Being under the care of caregivers (AOR=4.21; 95% CI: 1.31, 13.52), dressing fully during sun exposure (AOR=10.36; 95% CI: 3.37, 31.80), no oil massage during sun exposure (AOR=4.94; 95% CI: 2.01, 12.12), and being sick one month prior to the study (AOR=3.56; 95% CI: 1.51, 8.40) were positively associated with rickets.

Conclusion: One in every thirteen children had rickets in the study area. Being under the care of caregivers, dressing fully and no oil massage during sun exposure, and being sick one month prior to the study were significantly associated with rickets. Therefore, the health sectors should work to prevent rickets through counseling and encouraging of proper sunlight exposure and disease prevention.

Keywords: Eastern Ethiopia; Rickets; Under-five Children


How to cite: Sisay, K., Mesfin, F., Gobena, T. and Gebremichael, B. 2019. Rickets and Its Associated Factors among Under-Five Children in Selected Public Hospitals in Eastern Ethiopia. East African Journal of Health and Biomedical Sciences, Volume 3(2): 23-34.

Introduction

Rickets is a disease of children characterized by a failure or delay in endochondral calcification of the growth plates (physis) of long bones during times of rapid growth. It results in widening and splaying of the growth plates and leads to enlargement of the wrists and costochondral junctions, bowed legs and knock-knee (Pettifor and Prentice, 2011). There are various causes responsible for the different forms of rickets (Prentice, 2013), but it is primarily caused by the lack

of vitamin D, calcium and phosphate (Ahmed *et al.*, 2011).

Rickets is a universal problem and contributes to the high burden of illness and death among children below five years (Wondale *et al.*, 2005). It is reported worldwide (Matsuo *et al.*, 2009), including in the countries with a year-round opportunity for skin exposure to sunlight (Prentice, 2013). The problem is the most





¹Melka Kero Health Center, Dire Dawa, Ethiopia

²Federal Ministry of Health, Addis Ababa, Ethiopia

³Department of Environmental Health Sciences, College of Health and Medical Sciences, Haramaya University, Harar, Ethiopia

⁴School of Public Health, College of Health and Medical Sciences, Haramaya University, Harar, Ethiopia

common non-communicable diseases affecting children, particularly in developing countries (Hatun et al., 2005; Bishop, 2006; Bereket, 2010). It is most commonly seen in children from countries in Africa, the Middle East and Asia, although many of these countries are in the tropics with abundant sunshine (Craviari et al., 2008; Bener and Hoffmann, 2010; Aggarwal et al., 2012; Thacher et al., 2012; Prentice, 2013). The magnitude of rickets varies in different studies in the world, ranging from 3.4% in Kenya to 70.9% in Pakistan (Najada et al., 2004; Hatun et al., 2005; Bener and Hoffmann, 2010; Theuri, 2012; Uush, 2013; Khalil, 2014; Aziz et al., 2016; Sakr et al., 2016; Alshammari et al., 2018). In Ethiopia, the magnitude varies from 4% to 21.4% (Belachew et al., 2005b; Kenenisa et al., 2014; Derseh et al., 2018). There is one report from Ethiopia which found lack of exposure to sunshine and inadequate intake of vitamin D might be the cause for rickets (Wondale et al., 2005).

Rickets has the potential to produce an epidemic outbreak (Al-Atawi *et al.*, 2009), and if left untreated, it results in delay in developmental milestones, bone deformity and long term disability, pain, pathological fractures, obstructed labor, risk of falls and death (Misra *et al.*, 2008; Högler, 2015; Munns *et al.*, 2016), lower respiratory tract infections (Wondale *et al.*, 2005), restlessness, lack of sleep, underweight, and retarded growth (Nield *et al.*, 2006; Kenenisa *et al.*, 2014).

Many studies have shown that the development of rickets is related with low socioeconomic status (Agarwal et al., 2002; Wondale et al., 2005; Baroncelli et al., 2008; Manaseki-Holland et al., 2008; Bishay et al., 2016). In addition, being male (Bereket, 2006; Lips, 2007; Aziz et al., 2016), urban residence (Harinarayan et al., 2007; Lips, 2007), insufficient family income (Bishay et al., 2016), overcrowded living situation (Agarwal et al., 2002; Bereket, 2006; Grant et al., 2009), protein energy malnutrition (Wondale et al., 2005; Wagner and Greer, 2008), dark skin pigmentation (Bereket, 2006), reduced sunlight exposure (Bela-chew et al., 2005a; Bereket, 2006; Yassin and Lubbad, 2010; Kenenisa et al., 2014; Bishay et al., 2016; Mahmoud et al., 2016) and air pollution (Agarwal et al., 2002; Bereket, 2006) were significantly associated with the development of rickets.

Despite its burden and recommendations from many industrialized countries, there are no public health initiatives to address rickets prevention in resource-limited countries (Munns *et al.*, 2006; Mutlu *et al.*, 2011). Studies from some areas of Ethiopia indicate rickets is an important child health problem in the country (Belachew *et al.*, 2005b; Prentice, 2008; Pettifor, 2009; Kenenisa *et al.*, 2014; Derseh *et al.*, 2018). However, the magnitude and the associated factors of rickets in the eastern part of the country are not well studied and documented. Therefore, this study was aimed to assess the magnitude of rickets and the associated factors among under-five children in selected public hospitals of eastern Ethiopia.

Materials and Methods

Study design, area and period

A hospital based cross-sectional study was conducted among under-five children from February 23 to March 10, 2017 in selected public hospitals in eastern Ethiopia (Dilchora General Hospital, Haramaya General Hospital, and Hiwot Fana Specialized University Hospital).

Dilchora Hospital is one of the two public hospitals in Dire Dawa city, which is located at a distance of 515 kilometers to the east of Addis Ababa; capital city of Ethiopia. The hospital had 220 beds and 12 case teams to provide both inpatient and outpatient health care services to its catchment area. On the other hand, Haramaya Hospital is a general hospital, which is found in Haramaya town at a distance of 507 kilometers to the east of Addis Ababa. The hospital had 110 beds and 9 case teams to provide inpatient and outpatient health care services to the residents of Haramaya district and other neighboring districts. Hiwot Fana Hospital is a specialized University hospital which is found in Harar town at a distance of 526 kilometers to the East of Addis Ababa. The hospital had 163 beds and 12 case teams to provide referral inpatient and outpatient services to the residents of Harari Regional State and others surrounding it.

Sample size determination and sampling procedure

The sample size for this study was calculated by using a single population proportion formula considering the following assumptions: 95% confidence level, 2.5% margin of error, and the proportion of rickets among

under-five children from previous study in Jimma University Specialized Hospital, Southwestern Ethiopia (10.5%) (Kenenisa *et al.*, 2014). This resulted in sample size of 578. The final sample size was fixed at 636 after adding 10% for non-respondents.

The final sample size was allocated to each hospital proportionally based on number pediatrics patients attending each hospitals. The average number of pediatric patients flow for the past 15 days (i.e. data collection period of this study) was 450, 375 and 225 for Dilchora, Hiwot Fana and Haramaya Hospitals, respectively. Finally, all the children who visited the selected hospitals during the data collection were included in the study until the allocated sample size was attained.

Inclusion and exclusion criteria

Under-five children and their mothers/caregivers, who visited the Maternal and Child Health (MCH) clinic, pediatric Outpatient Department (OPD), nutrition rehabilitation unit and pediatrics inpatient ward of the selected public hospitals for health care services, were included. Seriously ill children and/or their mother/caregiver during the period of data collection were excluded from the study.

Data collection methods

Data were collected by the following methods;

Face-to-face interview: the children's mothers/care-givers were interviewed using semi-structured questionnaire by nine nurses who were supervised by three medical doctors. The questionnaire was developed by reviewing literatures, and it was used to collect data on socio-demographic and economic, associated factors, and dietary and non-dietary practices. A food frequency questionnaire was used to collect data on the consumption of different food groups during seven days before the study (CSA and WFP., 2019).

Anthropometric measurements: were done by the nurses who interviewed the mothers/caregivers. Digital weight scale was used to measure the weight of the children aged below two years, and their length was measured by wooden measuring board in lying position. For those who were two years old and above, digital weight scale, which had an attached height scale, was used to measure their weight and height. The weight scale was readjusted to zero and calibrated daily before data collection. During weight and height measurements, heavy clothes, shoes, bags or any other

material were avoided. Body weight was measured to the nearest 0.1 kg, and length/height was measured to the nearest 0.1 cm (Fryar *et al.*, 2012; Casadei and Kiel, 2019).

Clinical diagnosis of rickets: The clinical diagnosis of rickets was made by the joint decision of the data collectors (nurses) and the supervisors (medical doctors) using an instrument containing five clinical features which was used by various studies (Strand *et al.*, 2007; Uush, 2013; Edwards *et al.*, 2014). The clinical features include; age <5 years, length/height-for-age z-score <-2 Standard Deviation (SD), leg pain/ bow legs/knock-knees, wrist enlargement, and costochondral enlargement. The presence of at least three clinical features was used to diagnose rickets. The instrument had been reported of having sensitivity=87%, specificity=76%, positive predictive value=73%, and negative predictive value=89% (Thacher *et al.*, 2002).

Data quality control

To maintain the consistency of the data collection tool, the questionnaire was first prepared in English language and translated to Afan Oromo, Amharic and Somali languages, and back to English language by professional translators. An intensive training was provided to the data collectors and the supervisors for four days. In order to evaluate the acceptability and applicability of the procedures and tools, a pre-test was done on 32 (5%) of the sample in Sabian Hospital (Dire Dawa) one week before the actual data collection. To check the completeness and consistency of the questionnaire, the data collectors were closely supervised during the data collection process by the supervisors. Double data entry was done to minimize errors during data entry by checking the consistency between the two data sets.

Data analysis

The data were cleaned and entered into computer using EpiData Version 3.1 and exported into Statistical Package for the Social Sciences (SPSS) Version 22.0 for analysis. Categorical variables were described using frequency and percentage. Continuous variables were described using appropriate summary measures after they had been assessed for normality. To categorize the average family income, the Ethiopian national payroll and taxation guide was used (The Global Payroll Association, 2108).

Anthropometric indices with z-score values were computed using the World Health Organization (WHO) Anthro Version 3.2.2 in order to analyze the nutritional status of the children. A low height-for-age, below -2 SD of the reference population, indicates stunting while below -3 SD indicates severe stunting. A low weight-for-height, below -2 SD of the reference population, indicates wasting, while below -3 SD indicates severe wasting. A child with a weight-for-age z-score below -2 SD of the reference population is underweight while a child below -3 SD is severely underweight (WHO, 2006; Olack *et al.*, 2011).

In the bivariate analysis, Crude odds Ratio (COR) was computed to assess the association between each independent and the outcome variables. Then, the variables with p-value < 0.25 were included to multi-variable logistic regression analysis and the associated factors were identified by estimating Adjusted odds Ratio (AOR) with 95% Confidence Interval (CI). The final model was checked for goodness-of-fit using Hosmer-Lemeshow goodness-of-fit test. Finally, statistical significance was considered at p-value < 0.05.

Ethical considerations

Ethical clearance was secured from Institutional Health Research Ethics Review Committee (IHRERC) of the College of Health and Medical Sciences at Haramaya University. An official letter was written from Haramaya University to Dire Dawa City Administration Health Bureau, Hiwot Fana Specialized University Hospital and Haramaya District Health Office. Informed, voluntary, written and signed consent was obtained from each hospital's head, and child's parent/caregiver. The interviews and measurements were carried out privately in a separate room. Names and personal identifiers were not included in the questionnaire and checklist to ensure the confidentiality of participants' information. The mothers/ caregivers whose children had rickets were counseled by the data collectors and the supervisors. The children with acute under-nutrition were linked to the nutritional rehabilitation units of the respective hospitals.

Results

Socio-demographic characteristics of the study participants

Five hundred ninety under-five children with their mothers/caregivers were included in the final analysis

yielding a response rate of 92.8%. Three hundred two (51.2%) of the children were male. The mean (\pm SD) age of the children was 24.2 (±15.9) months and 132 (24.2%) were in the age group of 12-23 months. More than half, 333 (56.4%), of the children were residents of rural area. Three hundred eighty seven (65.6%) of the children were from Muslim families. Three hundred (51.5%) of the children were from households with family size of 5-7. Two hundred twelve (38.8%) of the children were from families with an average monthly income of 1651-3200 Ethiopian Birr (ETB). The mean $(\pm SD)$ age of the mothers/caregivers was $30.9 (\pm 9.0)$ years and 519 (88.0%) were married. Half of them (51.2%) were housewives. Regarding educational status of the mothers/caregivers, 286 (48.7%) of them were unable to read and write (Table 1).

Child feeding practice

Two-hundred eighty (47.5%) of the children were on breastfeeding during the study period; the frequency of feeding was 8-12 times per day for 93 (33.2%) of the children. The duration of breastfeeding was two years and above for 183 (59.0%) of the children who were not on breastfeeding during the study period. Many of the children (56.2%) started complementary feeding at the age of six months, and 297 (57.8%) were initiated with breast milk at birth. Most of the study participants (86.9%) reported that there was no recommendation of special foods for children, while 569 (96.4%) reported as there were no food restrictions in their communities such as uncooked/raw foods, meat, fish and egg (Table 2).

The results of this study revealed that milk/milk products and cereals/grains/tubers were the most frequently consumed food groups by the children. Accordingly, 391 (79.1%) and 290 (58.7%) of the children consumed milk/milk products and cereals/grains/tubers every day in the last seven days prior to the study, respectively. Fish/fish oil, 13 (2.6%), fruits, 19 (3.8%), and meat, 32 (6.5%), were the least frequently consumed food groups in the study area (Table 3).

Table 1: Socio-demographic characteristics of mothers/caregivers and their children who visited Hiwot Fana, Haramaya and Dilchora hospitals in eastern Ethiopia, 2017 (n=590).

Variables	Categories	Frequency	Percentage
Child birth order	First born	130	22.0
	Second born	159	26.9
	Third born	142	24.1
	Fourth and above	159	26.9
Child age (in completed months)	0-5	75	12.7
	6-11	88	14.9
	12-23	132	22.4
	24-35	128	21.7
	36-47	90	15.3
	48-59	77	13.1
Maternal/caregiver marital status	Married	519	88.0
	Single	30	5.1
	Divorced	20	3.4
	Widowed	21	3.5
Religion	Muslim	387	65.6
	Orthodox	141	23.9
	Protestant	62	10.5
Maternal/caregiver's occupation	Housewife	301	51.2
	Unemployed	26	4.4
	Self employed	73	12.4
	Government employed	86	14.6
	Farmer	102	17.3
Maternal//caregiver education	Unable to read and write	286	48.6
	Able to read and write	117	19.9
	Primary school(grade 1-8)	78	13.3
	Secondary school (grade 9-12)	37	6.3
	College and above	70	11.9
Family size (in number)	1-4	221	37.5
	5-7	304	51.5
	<u>></u> 8	65	11.0
Average monthly income (in ETB)* (n=547)	<u><</u> 600	20	3.7
	601-1650	124	22.7
	1651-3200	212	38.8
	3201-5250	127	23.2
	5251-7800	44	8.0
	7801-10900	20	3.7

^{*1}US\$ = 29.93 ETB

Table 2: Child feeding practices among mothers/caregivers of under-five children who visited Hiwot Fana, Ha-

ramaya and Dilchora hospitals in eastern Ethiopia, 2017.

Variable	Categories	Frequency	Percentage
Frequency of breastfeeding (n=280)	Per need	29	10.4
	<8 times	84	30.0
	8-12 times	93	33.2
	>12 times	74	26.4
Duration of breastfeeding (n=310)	Never	21	6.8
	6 months	31	10.0
	One year	75	24.2
	\geq 2 years	183	59.0
Time of initiation for complementary feeding (n=494)	Before six	113	22.9
	months		
	At six months	278	56.2
	After 6 months	103	20.9
First food given at birth (n=514)	Breast milk	297	57.8
-	Water	151	29.4
	Other foods*	66	12.8
Special foods recommended for children in the community	Yes**	77	13.1
(590)	No	513	86.9
Special foods restricted for children in the community (590)	Yes ***	21	3.6
	No	569	96.4

^{*}Butter, formula milk, honey, porridge, and fruits (juice); ** Milk, porridge and potato; *** uncooked/raw foods, meat, fish and egg

Non-dietary practices

Out of the 590 children included in this study, 556 (94.2%) of them were directly exposed to the sunlight. Among the children who were exposed to the sunlight, 232 (41.7%) were initiated exposure at the age of one month and above, and 277 (49.8%) were partially undressed during the exposure. Besides, 295 (53.1%) of the children were exposed openly to the sky, and 260 (46.8%) were exposed for 10-15 minutes. The common practice during the sun exposure was massaging the child with oil in 348 (62.6%) of the children. Four-

hundred sixty-seven (79.2%) of the children were found to be fully vaccinated for their age; while 191 (32.4%) were sick in the last month prior to the study (Table 4).

Nutritional status of children

Concerning the nutritional status of the children, 76 (12.9%) of them were found severely stunted. In addition, 130 (22.0%) were severely underweight. Furthermore, 175 (30.3%) of the children were severely wasted (Table 5).

Table 3: Consumption of different food groups (in frequency and percentage) in the seven days preceding the study among under-five children who visited Hiwot Fana, Haramaya and Dilchora hospitals in eastern Ethiopia, 2017 (n=494).

Variable	Frequency of consumption				_
	Never	< Once a week	1-2 times a week	3-6 times a week	Daily
Cereals/grains/tubers	52 (10.5%)	17 (3.4%)	11 (2.2%)	124 (25.1%)	290 (58.7%)
Legumes/pulses	102 (20.6%)	19 (3.8%)	22 (4.5%)	251 (50.8%)	100 (20.2%)
Fruits	266 (53.8%)	32 (6.5%)	99 (20.0%)	78 (15.8%)	19 (3.8%)
Vegetables	130 (26.3%)	115 (23.3%)	54 (10.9%)	80 (16.2%)	115 (23.3%)
Eggs	199 (40.3%)	145 (29.4%)	34 (6.9%)	45 (9.1%)	71 (14.4%)
Fish/fish oil	395 (80.0%)	73 (14.8%)	7 (1.4%)	6 (1.2%)	13 (2.6%)
Meat	296 (59.9%)	126 (25.5%)	14 (2.8%)	26 (5.3%)	32 (6.5%)
Milk/milk products	37 (7.5%)	24 (4.9%)	2 (0.4%)	40 (8.1%)	391 (79.1%)

Table 4: Non-dietary practices towards the prevention of rickets among under-five children who visited Hiwot Fana,

Haramaya and Dilchora hospitals eastern Ethiopia, 2017.

Variable	Categories	Frequency	Percentage	
Age at first sun exposure (n=556)	First week	61	11.0	
-	Second week	179	32.2	
	Third week	84	15.1	
	After one month	232	41.7	
Dressing during sun exposure (n=556)	Undressed	173	31.1	
	Partially undressed	277	49.8	
	Fully dressed	106	19.1	
Way of sun exposure (n=556)	Open to the sky	295	53.1	
	Under shade	156	28.1	
	Inside home	109	18.9	
Duration of sun exposure in minutes	<10	86	15.5	
(n=556)	10-15	260	46.8	
	15-20	169	30.4	
	>20	41	7.4	
Common practice during sun exposure	Oil massage	348	62.6	
(n=556)	Breast feeding	72	12.9	
	Nothing	136	24.5	
Child vaccinated for his/her age (590)	Yes	467	79.2	
	No	123	20.8	
Child sickness one moth prior to the	Yes*	191	32.4	
study (590)	No	399	67.6	

^{*}Pneumonia, upper respiratory tract infections, diarrhea and vomiting

Table 5: Nutritional status of under-five children who visited Hiwot Fana, Haramaya and Dilchora hospitals in eastern Ethiopia, 2017.

Nutritional indices	Nutritional status	Frequency	Percentage
Height-for-Age (z-score) (n=590)	Normal (≥ -2)	462	78.3
	Moderate stunting $(-3 \text{ to } < -2)$	52	8.8
	Severe stunting (< -3)	76	12.9
Weight-for-Age (z-score) (n=590)	Normal (\geq -2)	346	58.6
	Moderate underweight (-3 to < -2)	114	19.3
	Severe underweight (< -3)	130	22.0
Weight-for-Height (z-score) (n=577)	Normal (\geq -2)	262	45.4
	Moderate wasting (-3 to < -2)	140	24.3
	Severe wasting (< -3)	175	30.3

Magnitude of rickets and associated factors

The magnitude of rickets among the under-five children who visited the three hospitals was 46 (7.8%) (95% CI: 5.7, 10.2). In the bivariable analysis, late initiation of complementary feeding, full dressing during sun exposure, no oil massaging during sun exposure, being sick one month prior to the study, and underweight were the significantly associated factors with rickets among the under-five children (P<0.05).

In the multivariable analysis, being under the care of the caregiver, full dressing during sun exposure, no oil massaging during sun exposure, and being sick one month prior to the study were identified as the significantly associated factors of rickets among the underfive children (P<0.05).

The odds of rickets was 4.21 times higher among the children who were under the care of caregivers compared to those who were under the care of their mothers (AOR=4.21, 95% CI: 1.31, 13.52). Those children

who were exposed to sunlight with full dressing were 10.36 times more likely to have rickets compared to the undressed ones (AOR=10.36, 95% CI: 3.37, 31.80). The occurrence of rickets was 4.94 times higher in the children who were not on oil massaging during sun exposure compared to those who were on oil massage

(AOR=4.94, 95% CI: 2.01, 12.12). Moreover, the children who were sick in the last one month preceding the study were 3.56 times more likely to develop rickets compared to children who were not sick (AOR=3.56, 95% CI: 1.51, 8.40) (Table 6).

Table 6: Factors associated with rickets among under-five children who visited Hiwot Fana, Haramaya and Dilchora hospitals in eastern Ethiopia, 2017.

Variables	Categories	Ric	ckets	COR (95% CI)	AOR (95% CI) ^ф
		Yes (%)	No (%)	-	
Relation of respondent	Caregiver	8 (11.3)	63 (88.7)	1.61 (0.72, 3.60)	4.21 (1.31, 13.52)*
to the child	Mother	38 (7.3)	481 (92.7)	1	1
Age of mother/care-	≤ 30 years	32 (9.2)	315 (90.8)	1.66 (0.87, 3.19)	1.86 (0.75, 4.58)
giver	>30 years	14 (5.8)	229 (94.2)	1	1
Marital status of	Non-married	8 (11.3)	63 (88.7)	1.61 (0.72, 3.60)	1.53 (0.46, 5.08)
mother/caregiver	Married	38 (7.3)	481 (92.7)	1	1
Time for complemen-	< 6 month	9 (8.0)	104 (92.0)	1.33 (0.57, 3.08)	0.42 (0.13, 1.35)
tary feeding initiation	> 6 month	13 (12.6)	90 (87.4)	2.22 (1.04, 4.75)	1.78 (0.66, 4.77)
·	At six month	17 (6.1)	261 (93.9)	1	1
Recommended foods	Yes	9 (11.7)	68 (88.3)	1.70 (0.79, 3.68)	0.33 (0.12, 1.02)
for children in the community	No	37 (7.2)	476 (92.8)	1	1
Way of dressing during	Dressed fully	23 (21.7)	83 (78.3)	6.57 (2.71, 15.94)	10.36 (3.37, 31.80)**
sun exposure	Partially un- dressed	9 (3.2)	268 (96.8)	0.80 (0.29, 2.18)	0.81 (0.25, 2.62)
	Undressed	7 (4.0)	166 (96.0)	1	1
Common practice during sun exposure	No oil mas- sage	28 (13.5)	180 (86.5)	4.77 (2.32, 9.80)	4.94 (2.01, 12.12)**
	Oil massage	11 (3.2)	337 (96.8)	1	1
Immunization	No	13 (10.6)	110 (89.4)	1.55 (0.79, 3.05)	1.42 (0.52, 3.91)
	Yes	33 (7.1)	434 (92.9)	1	1
Sick in last one month	Yes	30 (15.7)	161 (84.3)	4.46 (2.37, 8.41)	3.56 (1.51, 8.40)*
	No	16 (4.0)	383 (96.0)	1	1
Underweight	Yes	26 (10.7)	218 (89.3)	1.94 (1.06, 3.57)	2.38 (0.99, 5.73)
	No	20 (5.8)	326 (94.2)	1	1

AOR: Adjusted Odds Ratio; CI: Confidence Interval; COR: Crude Odds Ratio; * final model goodness-of-fit using Hosmer-Lemeshow test (p=0.11); * significant at p<0.05; ** significant at p<0.001

Discussion

The magnitude of rickets was 7.8% in the study area. Being under the care of caregiver, full dressing during sun exposure, no oil massaging during sun exposure, and being sick in the last one month prior to the study were identified as positively associated factors of rickets among the under-five children.

The magnitude of rickets in this study is similar with the studies conducted in Ethiopia (10.5%) (Kenenisa *et al.*, 2014), Turkey (6%) (Hatun *et al.*, 2005), and Jordan (10.6%) (Najada *et al.*, 2004). However, it is higher than the magnitudes reported from Kenya

(3.4%) (Theuri, 2012) and Saudi Arabia (3.5%) (Khalil, 2014). The possible explanation for these variations may be due to differences in socio-economic, demographic and cultural characteristics, sample size, study period and diagnostic parameters.

Odds of rickets in our study was higher among the children under the care of caregivers than those under their mothers. This is consistent with the finding in Ethiopia (Belachew *et al.*, 2005b). This can be expressed as the affection between children and caregivers is not as strong as the affection between children and mothers, which results in poor child care like feeding and sunlight exposure.

One of the potential sources of vitamin D is synthesis in the skin from the ultraviolet-B light fraction of the sunlight. Decreased sunlight exposure occurs when sunlight is attenuated by clouds, air pollution, or shade. Lifestyles or cultural practices that decrease time spent outdoors or increase the amount of body surface area covered by clothing when outdoors further limit sunlight exposure. The effects of sunlight exposure on vitamin D synthesis are also decreased for individuals with darker skin pigmentation and by the use of sunscreens (Fuller and Casparian, 2001; Roth et al., 2005). The effect was seen in the current study; those children who were exposed to sunlight with full dressing were more likely to have rickets compared to fully undressed ones. This is supported by various previous studies (Belachew et al., 2005a; Belachew et al., 2005b; Bereket, 2006; Yassin and Lubbad, 2010; Kenenisa et al., 2014; Bishay et al., 2016; Mahmoud et al., 2016). However intense exposure to the solar ultraviolet rays during childhood and adolescence increases a person's risk of developing melanoma and basal cell carcinoma (Etzel et al., 2003).

The occurrence of rickets in this study was higher in children who were not on oil massage during sun exposure compared to their counterparts. There one report which indicates practice of oil massage and sunbath helps in the synthesis of vitamin D in the skin (Balasubramanian and Ganesh, 2008).

According to the current study, the odds of rickets was higher among the children who were sick in the last one-month preceding the study compared to their counterparts. This result is in line with study reports from Ethiopia (Wondale *et al.*, 2005) and Jordan (Najada *et al.*, 2004). This can be explained as rickets is a risk factor for infections particularly lower respiratory infection (Wondale *et al.*, 2005). This is because vitamin D may play a role in activating the innate immune system which the endogenous antimicrobial peptide can attracts monocytes and neutrophils and prevents viral and bacterial infections (Leikina *et al.*, 2005).

The study has some limitations; it might not show the temporal relation between the independent and dependent variables since it is a cross-sectional study. There is also the possibility that some of the responses might suffer from recall bias since the questions for

the dietary practices were based on recall. To minimize this bias, the data collectors were trained on how to probe the respondents by starting with the most recent food items consumed. In addition, the study participants were given sufficient time for adequate recall of long term memory. Also, sub-clinical rickets patients might have been missed since the diagnosis was based on clinical signs and symptoms. Furthermore, some children with other bone problems might have been diagnosed as having rickets as a result of similar signs and symptoms.

Conclusion

The study revealed that nearly one in thirteen children had rickets in the study area. Being under the care of the caregiver, full dressing during sun exposure, no oil massaging during sun exposure, and being sick in the last one month prior to the study were identified as positively associated factors of rickets among the under-five children. Therefore, the health sectors/health personals should work on the prevention of rickets through appropriate provision of information on sun exposure, oil messaging at time sun exposure and disease prevention. In addition, large scale studies supported by laboratory and radiological investigations should be conducted to produce more accurate magnitude estimates of rickets and its associated factors.

Acknowledgements

We would like to thank the administrative bodies and staffs of Hiwot Fana, Haramaya and Dilchora Hospitals for their cooperation. We would also like to thank the data collectors, supervisors, questionnaire translators, study participants and all who gave their hands in the study directly or indirectly.

Conflict of Interest

The authors declare that they have no competing interests.

Contribution of Authors

KS initiated the research, and all of the authors contributed in proposal writing, data entry and analysis, drafting and revising the paper, and manuscript writing.

References

Agarwal, K. S., Mughal, M. Z., Upadhyay, P., Berry, J. L., Mawer, E. B. and Puliyel, J. M. 2002. The impact

- of atmospheric pollution on vitamin D status of infants and toddlers in Delhi, India. *Archives of Disease in Childhood*, 87(2): 111–113.
- Aggarwal, V., Seth, A., Aneja, S., Sharma, B., Sonkar, P., Singh, S. 2012. Role of calcium deficiency in development of nutritional rickets in Indian children: a case control study. *Journal of Clinical medicine and Endocrinolgy and Metabolism*, 97(10): 3461–3466.
- Ahmed, S. F., Franey, C., McDevitt, H., Somerville, L., Butler, S., Galloway, P. 2011. Recent trends and clinical features of childhood vitamin D deficiency presenting to a children's hospital in Glasgow. *Archives of Disease in Childhood*, 96(7): 694–696.
- Al-Atawi, M. S., Al-Alwan, I. A., Al-Mutair, A. N., Tamim, H. M. and Al-Jurayyan, N. A. 2009. Epidemiology of Nutritional Rickets in Children. Saudi Journal of Kidney Disease and Transplantation, 20(2): 260-265.
- Alshammari, O. M. O., Almezani, A. M. M., Alshlaqy, A. S., Alsiraa, M. N. N., Alenazy, G. A. G. and Ghamdi, S. A. S. K. 2018. The Prevalence of Rickets Disorder among Children in Saudi Arabia. *The Egyptian Journal of Hospital Medicine*, 73(6): 6943-6947.
- Aziz, K. T., Lehrasab, W., Ahmed, N. and Batool, Z. 2016.
 Frequency of Nutritional Rickets in Children 259 Months of Age with Severe Pneumonia.
 Annals of Pakistan Institute of Medical Sciences,
 12(2): 108-112.
- Balasubramanian, S. and Ganesh, R. 2008. Vitamin D deficiency in exclusively breast-fed infants. *Indian Journal of Medical Research*, 127(2008): 250-255.
- Baroncelli, G. I., Bereket, A., Kholy, M. E., Audì, L., Cesur, Y., Ozkan, B. 2008. Rickets in the Middle East: Role of environment and genetic predisposition. *The Journal of Clinical Endocrinology and Metabolism*, 93(5): 1743-1750.
- Belachew, T., Gebremariam, A., Legesse, W., Asres, T., Tiku, S., Kebede, E. 2005a. Module on micronutrient deficiency for the Ethiopian health center team. Jimma University in collaboration with Ethiopia Public Health Training Initiative, the Carter Center, the Ethiopia Ministry of Health, and the Ethiopia Ministry of Education. https://www.cartercenter.org/resources/pdfs/hea

- lth/ephti/library/modules/FinalModuleMicronut rientDeficiency.pdf[Accessed November 21, 2016]
- Belachew, T., Nida, H., Getaneh, T., Woldemariam, D. and Getinet, W. 2005b. Calcium Deficiency and Causation of Rickets in Ethiopian Children. *East African Medical Journal*, 82(3): 154-160
- Bener, A. and Hoffmann, G. F. 2010. Nutritional rickets among children in a sun rich Country. *International Journal of Pediatric Endocrinology*, 410502 (2010): doi:10.1155/2010/410502.
- Bereket, A. 2006. Nutritional rickets: nature or nurture?. Expert Review of Endocrinology & Metabolism, 1(5): 661–671.
- Bereket, A. 2010. Nutritional rickets: still a problem for the pediatric population. *Pediatric Health*, 4(1):75–87
- Bishay, S. N., El-Sherbini, M. H., Azzam, A. A. and Lotfy, A. A. 2016. Incidence and Risk Factors of Rachitic Genu Varus in Preschool Children in a Paediatric Health Institute in Egypt as One of the Developing Countries. *Open Orthopaedics Journal*, 10 (2016): 412-419. doi: 10.2174/1874325001610010412.
- Bishop, N. 2006. Don't ignore vitamin D. *Archives of Disease in Childhood*, 91(7): 549–550.
- Casadei, K. and Kiel, J. 2019. Anthropometric measurement. [Updated 2019 Mar 24]. In: StatPearls [Internet]. Treasure Island (FL): Stat PearlsPublishing. https://www.ncbi.nlm.nih.gov/books/NBK53

7315/ [Accessed April 10, 2019]

- Craviari, T., Pettifor, J. M., Thacher, T. D., Meisner, C., Arnaud, J., Fischer, P. R. 2008. Rickets: an overview and future directions, with special reference to Bangladesh. A summary of the Rickets Convergence Group meeting, Dhaka, 26-27 January 2006. *Journal of Health, Population Nutrition*, 26(1):112–121.
- CSA and WFP. 2019. Comprehensive Food Security and Vulnerability Analysis (CFSVA) of Ethiopia. https://reliefweb.int/sites/reliefweb.int/files/reso urces/wfp_ethiopia_cfsva_report_june_2019.pd f [Accessed June 05, 2019].
- Derseh, B., Mruts, K., Demie, T. and Gebremariam, T. 2018. Co-morbidity, treatment outcomes and factors affecting the recovery rate of under -five children with severe acute malnutrition admitted in selected hospitals from Ethiopia: retrospective

- follow up study. *Nutrition Journal*, 17:116.https://doi.org/10.1186/s12937-018-0423-1.
- Edwards, J. K., Thiongó, A., Van den Bergh, R., Kizito, W., Kosgei, R. J., Sobry. 2014. Preventable but neglected: rickets in an informal settlement, Nairobi, Kenya. *Public Health Action*, 4(2): 122–127.
- Etzel, R. A., Balk, S. J. and editors. 2003. How Environmental Exposures Influence the Development and Exacerbation of Asthma. *American Academy of Pediatrics*, 112 (Suppl. 1):233-239.
- Fryar, C. D., Gu, Q. and Ogden, C. L. 2012. Anthropometric reference data for children and adults: United States, 2007–2010. National Center for Health Statistics. Vital Health Stat, 11(252): 2012-1602.
- Fuller, K. E. and Casparian, J. M. 2001. Vitamin D: balancing cutaneous and systemic considerations. *Southern Medical Journal*, 94(1):58-64
- Grant, C. C., Wall, C. R., Crengle, S. and Scragg, R. 2009. Vitamin D deficiency in early childhood: prevalent in the sunny South Pacific. *Public Health Nutrition*, 12(10): 1893-1901.
- Harinarayan, C. V., Ramalakshmi, T., Prasad, U. V., Sudhakar, D., Srinivasarao, P. V. and Kadainti, V. S. 2007. High prevalence of low dietary calcium, high phytate consumption, and vitamin D deficiency in healthy South Indians. *Journal of Clinical Nutrition*, 85(4): 1062-1067.
- Hatun, S., Ozkan, B., Orbak, Z., Doneray, H.,
 Cizmecioglu, F., Toprak, D. 2005. Vitamin
 D Deficiency in Early Infancy. *Journal of Nutrition*, 135(2): 279–282.
- Högler, W. 2015. Complications of vitamin D deficiency from the foetus to the infant: One cause, one prevention, but who's responsibility? . Best Practice & Research Clinical Endocrinology & Metabolism, 29(3): 385-398.
- Kenenisa, C., Ewnetu, H. and Sime, H. 2014.
 Retrospective Analysis of Prevalence of Rickets and Associated Factors among Children Admitted to Pediatric Ward in Jimma University Specialized Hospital. *Journal of Pediatrics and Neonatal Care Pediatr Neonatal Care*, 1(7):00044.
- Khalil, T. 2014. Prevalence of rickets among children below one-year encounter of North West Armed

- Forced Hospital in Tabuk. *International Journal of Medical Sciences and Public Health*, 3(7):827-831.
- Leikina, E., Delanoe-Ayari, H., Melikov, K., Cho, M.S., Chen, A., Waring, A.J. 2005. Carbohydrate-binding molecules inhibit viral fusion and entry by crosslinking membrane glycoproteins. *Nature Immunology*, 6(10): 995-100.
- Lips, P. 2007. Vitamin D status and nutrition in Europe and Asia. *Journal of Steroid Biochemistry and Molecular Biology*, 103(3-5): 620-625.
- Mahmoud, A. O., Ahmed, A. Y. and Aly, H. M. 2016. The prevalence of active nutritional rickets in Egyptian infants in Cairo. *Egyptian Pediatric Association Gazette*, 64(3):105–110.
- Manaseki-Holland, S., Zulf, M. M., Bhutta, Z. and Qasem, S. M. 2008. Vitamin D status of socio-economically deprived children in Kabul, Afghanistan. *International Journal for Vitamin and Nutrition Research*, 78(1): 16-20.
- Matsuo, K., Mukai, T., Suzuki, S. and Fujieda, K. 2009. Prevalence and risk factors of vitamin D deficiency rickets in Hokkaido. *Pediatrics International*, 51 (2009): 559–562.
- Misra, M., Pacaud, D., Petryk, A., Collett-Solberg, P. F. and Kappy, M. 2008. Vitamin D deficiency in children and its management: review of current knowledge and recommendations. *Pediatrics*, 122 (2) 398-417.
- Munns, C., Zacharin, M. R., Rodda, C. P., Batch, J. A., Morley, R., Cranswick, N. E. 2006. Prevention and treatment of infant and childhood vitamin D deficiency in Australia and New Zealand: a consensus statement. *Medical Journal of Australia*, 185(5):268-72.
- Munns, C. F., Shaw, N., Kiely, M., Specker, B. L., Thacher, T. D. and Ozono, K. 2016. Global Consensus Recommendations on Prevention and Management of Nutritional Rickets. *Journal of Clinical Endocrinology and Metabolism*, 101(2): 394–415.
- Mutlu, G. Y., Kusdal, Y., Ozsu, E., Cizmecioglu, F. M. and Hatun, S. 2011. Prevention of Vitamin D deficiency in infancy. *International Journal of Pediatrics and Endocrinology*, 2011(1):4. doi: 10.1186/1687-9856-2011-4.

- Najada, A. S., Habashneh. M.S. and M., K. 2004. The frequency of nutritional rickets among hospitalized infants and its relation to respiratory diseases. *Journal of Tropical Pediatrics*, 50(6): 364-368.
- Nield, L. S., Mahajan, P., Joshi, A. and Kamat, D. 2006. Rickets: Not a disease of the past. American Family Physician, 74(4): 619-626.
- Olack, B., Burke, H., Cosmas, L., Bamrah, S., Dooling, K., Feikin, D. R. 2011. Nutritional status of under-five children living in an informal urban settlement in Nairobi, Kenya. *Journal of Health, Population Nutrition*, 29(4): 357-363.
- Pettifor, J. 2009. Vitamin D and calcium nutrition in children in developing countries. *Bull NutrFnd India*, 31 (2): 607-610.
- Pettifor, J. M. and Prentice, A. 2011. The role of vitamin D in paediatric bone health. Clinical Endocrinology and Metabolism, 25(4): 573–584.
- Prentice, A. 2008. Vitamin D deficiency: a global perspective. *Nutrition Reviews*, 66 (suppl 2): S153–S164.
- Prentice, A. 2013. Nutritional rickets around the world. *Journal of Steroid Biochemistry and Molecular Biology*, 136(2013): 201–206.
- Roth, D. E., Martz, P., Yeo, R., Prosser, C., Bell, M. and Jones, A. B. 2005. Are national vitamin D guidelines sufficient to maintain adequate blood levels in children? *Canadian Journal of Public Health*, 96 (6): 443-449.
- Sakr, M. A., Abdel-Aal, M. I., Zannoun, M. A. and Al-Maaty, M. M. A. 2016. Prevalence and clinical characteristics of rickets in infants ans children attending the outpatient clinic at Damietta University Hospital. *Asian Journal of Clinical Nutrition*, 8 (1-3): 9-13
- Strand, M. A., Perry, J., Jin, M., Tracer, D. P., Fischer, P. R., Zhang, P. 2007. Diagnosis of rickets and reassessment of prevalence among rural children in northern China.

- Pediatrics International, 49(2007): 202–209.
- Thacher, T. D., Fischer, P. R., Isichei, C. O., Zoakah, A. I. and Pettifor, J. M. 2012. Prevention of nutritional rickets in Nigerian children with dietary calcium supplementation. *Bone*, 50 (5): 1074–1080.
- Thacher, T. D., Fischer, P. R. and Pettifor, J. M. 2002. The usefulness of clinical features to identify active rickets. *Annals of Tropical Paediatrics*, 22(3): 229–237.
- The Global Payroll Association. 2108. Ethiopia payroll and tax overview: a guide to doing business in Ethiopia.https://cdn.shopify.com/s/files/1/2358/5863/files/Ethiopia_2018_v2.pdf [Accessed September 16, 2019]
- Theuri, A. W. 2012. Factors associated with rickets among children aged 0-59 months at Kiambu District Hospital, Kenya. *University of Nairobi researcharchive*:6–22. http://erepository.uonbi.ac.ke/bitstream/hand le/11295/6793/Abstract.pdf?sequence=1 [Accessed October 09, 2016].
- Uush, T. 2013. Prevalence of classic signs and symptoms of rickets and vitamin D deficiency in Mongolian children and women. *The Journal of Steroid Biochemistry and Molecular Biology*, 136(2013): 207–210.
- Wagner, C. L. and Greer, F. R. 2008. Prevention of rickets and vitamin D deficiency in infants, children and adolescents. *Pediatrics*, 122 (5) 1142-1152.
- WHO. 2006. WHO child growth standards: length/height-for-age, weight-for-length, weight-for-height and body mass index-for-age. Geneva: World Health Organization.
- Wondale, Y., Shiferaw, F. and Lulseged, S. 2005. A systematic review of nutritional rickets in Ethiopia: status and prospects. *Ethiopian Medical Journal*, 43(3): 203-210.
- Yassin, M. M. and Lubbad, A. M. H. 2010. Risk factors associated with nutritional rickets among children aged 2 to 36 months old in the gaza strip: a case control study. *International Journal of Food, Nutrition and Public Health*, 3(1): 45-33.